

REGULAR GRADED ALGEBRAS AND OBSTRUCTED CATEGORIES WITH DUALITY

STEVEN DUPLIJ AND WLADYSLAW MARCINEK

ABSTRACT. Regular and higher regular graded algebras (in simplest case satisfying Von Neumann regularity $\Theta_1\Theta_2\Theta_1 = \Theta_1$ instead of anticommutativity) are introduced and their properties are studied. They are described in terms of obstructed categories with nonclosed invertible and noninvertible morphisms for which generalized (obstructed) functors and natural transformations are given. Regular algebras and bialgebras are considered with examples. Corresponding regularization of the cross product and Wick algebras is made.

1. INTRODUCTION

The concept of higher regularization was introduced in the abstract way in the supermanifold theory [9, 8] and then considered for general morphisms [12, 11], which led to regularization of categories and Yang-Baxter equation [13, 14, 15]. This concept is close to the some generalizations of category theory [1, 3, 4, 16], it is connected with weak bialgebras [24, 25], squared Hopf algebras [21] and weak Hopf algebras [19, 10, 20] and plays an important role in topological quantum field theories [7, 2]. Here we introduce higher regular graded algebras and study them in terms of obstructed categories, which can clear and concrete understanding of the concept of semisupermanifold [9].

Let $\mathcal{A} = \mathcal{A}_{\overline{0}} \oplus \mathcal{A}_{\overline{1}}$ be an arbitrary superalgebra equipped with an associative multiplication $m : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$ [16]. We use the notation $m(a \otimes b) \equiv ab$ for the multiplication (we call it internal product). Denote by $|a|$ the parity of $a \in \mathcal{A}$. Then we have the relation $|ab| = |a| + |b| \pmod{2}$ for the multiplication in $\mathcal{A}$. The multiplication m is graded homogeneous mapping

$$(1) \quad m_{\overline{i}\overline{j}} : \mathcal{A}_{\overline{i}} \otimes \mathcal{A}_{\overline{j}} \rightarrow \mathcal{A}_{\overline{i}+\overline{j} \pmod{2}},$$

where $m_{\overline{i}\overline{j}}$ is the restriction of m to $\mathcal{A}_{\overline{i}} \otimes \mathcal{A}_{\overline{j}}$ and $\overline{i}, \overline{j} = 0, 1$. It is obvious that the mapping $m_{\overline{0}\overline{0}} : \mathcal{A}_{\overline{0}} \otimes \mathcal{A}_{\overline{0}} \rightarrow \mathcal{A}_{\overline{0}}$ is a multiplication turning $\mathcal{A}_{\overline{0}}$ into a closed associative algebra. Mappings $m_{\overline{0}\overline{1}}$ and $m_{\overline{1}\overline{0}}$ define on $\mathcal{A}_{\overline{1}}$ an $\mathcal{A}_{\overline{0}}$ -module structure. The mapping $m_{\overline{1}\overline{1}} : \mathcal{A}_{\overline{1}} \otimes \mathcal{A}_{\overline{1}} \rightarrow \mathcal{A}_{\overline{0}}$ is an $\mathcal{A}_{\overline{0}}$ -valued bilinear form on $\mathcal{A}_{\overline{1}}$. This means that the product ab is an even element of $\mathcal{A}$, $|ab| = 0 \pmod{2}$ for arbitrary two odd elements $a, b \in \mathcal{A}_{\overline{1}}$. Supercommutativity [5] means that $ab - (-1)^{|a||b|}ba = 0$. Let us consider a graded algebra $\mathcal{A}$ generated by a set $E \subset \mathcal{A}_{\overline{1}}$ of odd elements. In this case $\mathcal{A}_{\overline{0}}$ is the image of $m_{\overline{1}\overline{1}}$

$$(2) \quad \mathcal{A}_{\overline{0}} := \text{Im}(m_{\overline{1}\overline{1}}).$$

Note that $\mathcal{A}_{\overline{1}}$ is also closed under the triple product $m_{\overline{1}\overline{1}\overline{1}} : \mathcal{A}_{\overline{1}} \otimes \mathcal{A}_{\overline{1}} \otimes \mathcal{A}_{\overline{1}} \rightarrow \mathcal{A}_{\overline{1}}$, where

$$(3) \quad m_{\overline{1}\overline{1}\overline{1}} := m_{\overline{0}\overline{1}} \circ (m_{\overline{1}\overline{1}} \times \text{id}) = m_{\overline{1}\overline{0}} \circ (\text{id} \times m_{\overline{1}\overline{1}}).$$

2. REGULAR GRADED ALGEBRAS

Here we consider graded algebras $\mathcal{A}^{reg}$ as a direct product of superalgebras with new multiplication $(\cdot)$ (we call it external product) between copies satisfying the following regularity relations

$$(4) \quad \begin{aligned} \mathbf{a}_1 \cdot \mathbf{a}_2 \cdot \mathbf{a}_1 &= \mathbf{a}_1, & \mathbf{a}_2 \cdot \mathbf{a}_1 \cdot \mathbf{a}_2 &= \mathbf{a}_2, \\ \mathbf{a}_1 &\in \mathcal{A}_1, & \mathbf{a}_2 &\in \mathcal{A}_2, & \mathcal{A}^{reg} &= \mathcal{A}_1 \times \mathcal{A}_2, \end{aligned}$$

or their higher order generalizations [12, 11]. We consider a class of graded algebras as a free product of n copies of a superalgebra $\mathcal{A}$ modulo the regularity relations (4) and call them *regular graded algebras*.

Recall ([6] and refs. therein) that the free product of algebras $\mathcal{A}_1$ and $\mathcal{A}_2$ is the algebra $\mathcal{A}_1 \times \mathcal{A}_2$ formed by all formal finite sums of monomials of the form $\mathbf{a}_1 \cdot \mathbf{b}_2 \cdot \mathbf{a}_1 \dots$ or $\mathbf{b}_1 \cdot \mathbf{a}_2 \cdot \mathbf{b}_1 \dots$, where $\mathbf{a}_i, \mathbf{b}_i \in \mathcal{A}_i$, $i = 1, 2$, without relations between them.

Let us consider higher generalizations of (4).

Definition 1. An element $\mathbf{a}_1 \in \mathcal{A}_1$ for which there are elements $\mathbf{a}_i \in \mathcal{A}_i$, where $i = 2, \dots, n$ satisfying the following condition

$$(5) \quad \mathbf{a}_1 \cdot \mathbf{a}_2 \cdot \dots \cdot \mathbf{a}_n \cdot \mathbf{a}_1 = \mathbf{a}_1,$$

and its all cyclic permutations is said to be *n-regular* in $\mathcal{A} = \mathcal{A}_1 \times \mathcal{A}_2 \times \dots \times \mathcal{A}_n$. The set of all *n-regular* elements of $\mathcal{A}$ is denoted by $\mathcal{A}^{reg} = Reg^{(n)}(\mathcal{A})$.

The above *n-regularity* condition is essential for all noninvertible elements of the algebra $\mathcal{A}$.

Definition 2. If the product of arbitrary two *n-regular* elements $\mathbf{a}, \mathbf{b} \in Reg^{(n)}(\mathcal{A})$ is also *n-regular* $\mathbf{a} \cdot \mathbf{b} \in Reg^{(n)}(\mathcal{A})$, then the multiplication $m^{reg}(\mathbf{a} \otimes \mathbf{b}) = \mathbf{a} \cdot \mathbf{b}$ is said to be *n-regular*.

The (super-)algebra $\mathcal{A}$ generating by *n-regular* elements and equipped with *n-regular* multiplication is said to be *n-regular*.

3. REGULAR ODD GRADED ALGEBRAS

As a clear example we consider graded algebras $\mathcal{A} = \mathcal{A}_{\overline{0}} \oplus \mathcal{A}_{\overline{1}}$ generated by *n-regular* odd elements $Reg(\mathcal{A}_{\overline{1}})$ which we call *regular odd graded algebras*. Obviously, that for these algebras every monomial formed by an external product of an even number of generators is an even element of $\mathcal{A}$, and the monomial with an odd number of generators is an odd element. The superalgebra $\mathcal{A}$ contains all $\mathbb{K}$ -linear combinations of such monomials and external products.

We consider as an example an associative graded algebra $\mathcal{A} = \mathcal{A}^{reg}(\Theta_1, \Theta_2)$ generated by two noncommuting odd generators Θ_1 and Θ_2 satisfying the following relations

$$(6) \quad \Theta_1^2 = \Theta_2^2 = 0.$$

and

$$(7) \quad \Theta_1 \Theta_2 \Theta_1 = \Theta_1, \quad \Theta_2 \Theta_1 \Theta_2 = \Theta_2.$$

So the graded algebra $\mathcal{A} = \mathcal{A}^{reg}(\Theta_1, \Theta_2)$ is regular, but *not* anticommutative. Let us introduce

$$E_{12} = \Theta_1 \Theta_2, \quad E_{21} = \Theta_2 \Theta_1.$$

It is easy to observe that E_{12} and E_{21} are idempotents, since

$$\begin{aligned} E_{12}E_{12} &= (\Theta_1\Theta_2\Theta_1)\Theta_2 = \Theta_1\Theta_2 = E_{12}, \\ E_{21}E_{21} &= (\Theta_2\Theta_1\Theta_2)\Theta_1 = \Theta_2\Theta_1 = E_{21}. \end{aligned}$$

Note that idempotents E_{ij} are orthogonal $E_{12}E_{21} = E_{21}E_{12} = 0$ and associative, and therefore they form an idempotent semigroup [26, 27]. From (7) it follows that E_{ij} are two-sided units for Θ_i , viz. $E_{12}\Theta_1 = \Theta_1$, $\Theta_1E_{21} = \Theta_1$, $E_{21}\Theta_2 = \Theta_2$, $\Theta_2E_{12} = \Theta_2$. A generic element in $\mathcal{A}^{reg}(\Theta_1, \Theta_2)$ is

$$(8) \quad \mathbf{a} = a_0 + a_1\Theta_1 + a_2\Theta_2 + a_{12}\Theta_1\Theta_2 + a_{21}\Theta_2\Theta_1$$

The multiplication in $\mathcal{A}^{reg}(\Theta_1, \Theta_2)$ is given by

$$(9) \quad \begin{aligned} \mathbf{a}\mathbf{b} &= a_0b_0 \\ &+ (a_0b_1 + a_1b_0 + \underline{a_1b_{21}} + \underline{a_{12}b_1})\Theta_1 \\ &+ (a_0b_2 + a_2b_0 + \underline{a_2b_{12}} + \underline{a_{21}b_2})\Theta_2 \\ &+ (a_0b_{12} + a_1b_2 + a_{12}b_0 + \underline{a_{12}b_{12}})\Theta_1\Theta_2 \\ &+ (a_0b_{21} + a_2b_1 + a_{21}b_0 + \underline{a_{21}b_{21}})\Theta_2\Theta_1. \end{aligned}$$

Note that the underlined terms are absent in anticommutative algebras [5], and so they appear *only* due to regularity (7). That considerably changes the whole machinery of regular graded algebras computations as compared with anticommutative case [5]. For instance, the equation $\mathbf{a}\mathbf{b} = 0$ has the solution $\mathbf{b}_{zdiv} = b_0(1 - \Theta_1 - \Theta_2)$, which means that elements $\mathbf{b}_{zdiv}$ are zero divisors for all elements of the algebra $\mathcal{A}^{reg}(\Theta_1, \Theta_2)$. For comparison, in $\Lambda(\theta_1, \theta_2)$ (see e.g. [5]) we have $\mathbf{b}_{zdiv}^\Lambda = k(a_2\theta_1 + a_1\theta_2) + r\theta_1\theta_2$, where k and r are any in a given field. Then, instead of well known formula for inverse in $\Lambda(\theta_1, \theta_2)$ as ($a_0 \neq 0$)

$$(a_0 + a_1\theta_1 + a_2\theta_2 + a_{12}\theta_1\theta_2)^{-1} = a_0^{-1} - a_0^{-2}(a_1\theta_1 + a_2\theta_2 + a_{12}\theta_1\theta_2)$$

we have in $\mathcal{A}^{reg}(\Theta_1, \Theta_2)$

$$\begin{aligned} \mathbf{a}^{-1} &= a_0^{-1} - D^{-1}[a_1\Theta_1 + a_2\Theta_2 \\ &+ \left(a_{12}\left(1 + \frac{a_{21}}{a_0}\right) - \frac{a_1a_2}{a_0}\right)\Theta_1\Theta_2 + \left(a_{21}\left(1 + \frac{a_{12}}{a_0}\right) - \frac{a_1a_2}{a_0}\right)\Theta_2\Theta_1], \end{aligned}$$

where $D = (a_0 + a_{12})(a_0 + a_{21}) - a_1a_2$.

4. REGULAR GRADED ALGEBRA REPRESENTATIONS

The graded algebra $\mathcal{A} = \mathcal{A}^{reg}(\Theta_1, \Theta_2)$ can be presented by a free product of two one dimensional Grassmann algebras $\Lambda(\theta_1)$ and $\Lambda(\theta_2)$ modulo regularity relations (4) (denote them by $\overset{reg}{\sim}$) as

$$\mathcal{A}^{reg}(\Theta_1, \Theta_2) \cong \Lambda(\theta_1) \times \Lambda(\theta_2) / \overset{reg}{\sim}.$$

That means $\Theta_1\Theta_2\Theta_1 = \Theta_1 \iff \theta_1 \cdot \theta_2 \cdot \theta_1 = \theta_1$, where the external $(\cdot)$ -product of elements from different copies of one dimensional Grassmann algebra is defined in (4).

Let us consider n copies of one-dimensional Grassmann algebra $\Lambda(\theta_1)$. The i -th copy we denote by $\Lambda(\theta_n)$. Let us define a graded algebra $\mathcal{A}^{reg}(\Theta_1, \Theta_2, \dots, \Theta_n)$ as a free product of n copies of one-dimensional Grassmann algebras subject to the following relation

$$(10) \quad \Theta_1\Theta_2\Theta_3 \dots \Theta_n\Theta_1 = \Theta_1$$

and its all cyclic permutations. So the graded algebra $\mathcal{A}^{reg}(\Theta_1, \Theta_2, \dots, \Theta_n)$ is regular. Thus the graded algebra $\mathcal{A} = \mathcal{A}^{reg}(\Theta_1, \Theta_2, \dots, \Theta_n)$ can also be presented by a free product of n one dimensional Grassmann algebras $\Lambda(\theta_1)$ modulo regularity relations (4) (denote them by $\overset{reg}{\sim}$) as

$$\mathcal{A}^{reg}(\Theta_1, \Theta_2, \dots, \Theta_n) \cong \Lambda(\theta_1) \times \Lambda(\theta_2) \times \dots \times \Lambda(\theta_n) / \overset{reg}{\sim}.$$

For the n -regular graded algebra $\mathcal{A}^{reg}(\Theta_1, \Theta_2, \dots, \Theta_n)$ we introduce a collection of right and left multiplication operators R_{Θ_i} and L_{Θ_i} by the formulas

$$(11) \quad R_{\Theta_i}(x) := x\Theta_i, \quad L_{\Theta_i}(x) := \Theta_i x,$$

for every $x \in \mathcal{A}^{reg}(\Theta_1, \Theta_2, \dots, \Theta_n)$ and $i = 1, \dots, n$.

Lemma 3. *The correspondence defines a representation of $\mathcal{A}^{reg}(\Theta_1, \Theta_2, \dots, \Theta_n)$ into itself*

$$(12) \quad \mathcal{L} : \Theta_i \rightarrow L_{\Theta_i} \in \text{End}(\mathcal{A}^{reg}(\Theta_1, \Theta_2, \dots, \Theta_n))$$

and

$$(13) \quad \mathcal{R} : \Theta_i \rightarrow R_{\Theta_i} \in \text{End}(\mathcal{A}^{reg}(\Theta_1, \Theta_2, \dots, \Theta_n))$$

defines an antirepresentation of $\mathcal{A}^{reg}(\Theta_1, \Theta_2, \dots, \Theta_n)$ into itself.

Proof: We calculate

$$\begin{aligned} R_{\Theta_i} R_{\Theta_j} &= R_{\Theta_i}(R_{\Theta_j}(x)) = R_{\Theta_i}(x\Theta_j) = x\Theta_j\Theta_i = R_{\Theta_j\Theta_i}, \\ L_{\Theta_i} L_{\Theta_j} &= L_{\Theta_i}(L_{\Theta_j}(x)) = L_{\Theta_i}(\Theta_j x) = \Theta_i\Theta_j x = L_{\Theta_i\Theta_j}. \end{aligned}$$

5. REGULAR GRADED ALGEBRA DECOMPOSITION

Now we use Lemma 3 to construct the graded algebra decomposition into linear spaces. First we consider the regular graded algebra $\mathcal{A}^{reg}(\Theta_1, \Theta_2)$ and introduce 2 two-dimensional linear spaces $\mathbf{X}_1$ and $\mathbf{X}_2$ over a field $\mathbb{K}$

$$(14) \quad \begin{aligned} \mathbf{X}_1 &:= \text{lin}_{\mathbb{K}}\{\Theta_1, \Theta_1\Theta_2\} = \text{lin}_{\mathbb{K}}\{\Theta_1, E_{12}\}, \\ \mathbf{X}_2 &:= \text{lin}_{\mathbb{K}}\{\Theta_2, \Theta_2\Theta_1\} = \text{lin}_{\mathbb{K}}\{\Theta_2, E_{21}\}. \end{aligned}$$

Define two linear mappings $f_1 : \mathbf{X}_1 \rightarrow \mathbf{X}_2^T$ and $f_2 : \mathbf{X}_2 \rightarrow \mathbf{X}_1^T$ (where T is coordinate transposition, $T^2 = 1$) as a left multiplication by Θ_2 and Θ_1 respectively (because $\mathcal{L}$ is a representation). We obtain

$$(15) \quad \begin{aligned} f_1(\Theta_1) &= L_{\Theta_2}(\Theta_1) = \Theta_2\Theta_1, & f_1(\Theta_1\Theta_2) &= L_{\Theta_2}(\Theta_1\Theta_2) = \Theta_2\Theta_1\Theta_2 = \Theta_2, \\ f_2(\Theta_2) &= L_{\Theta_1}(\Theta_2) = \Theta_1\Theta_2, & f_2(\Theta_2\Theta_1) &= L_{\Theta_1}(\Theta_2\Theta_1) = \Theta_1\Theta_2\Theta_1 = \Theta_1, \end{aligned}$$

where regularity plays an essential role. Obviously we have

$$(16) \quad f_1 \circ f_2 \circ f_1 = f_1, \quad f_2 \circ f_1 \circ f_2 = f_2.$$

Define two mappings

$$(17) \quad e_{\mathbf{X}_1} := f_2 \circ f_1, \quad e_{\mathbf{X}_2} := f_1 \circ f_2.$$

It is obvious that we have

$$(18) \quad e_{\mathbf{X}_1} \circ e_{\mathbf{X}_1} = e_{\mathbf{X}_1}, \quad e_{\mathbf{X}_2} \circ e_{\mathbf{X}_2} = e_{\mathbf{X}_2}.$$

Observe that mappings $e_{\mathbf{X}_i}$ are identities only for regular graded algebras $\mathcal{A}^{reg}(\Theta_1, \Theta_2)$. Thus we can construct a category $\mathcal{C}_{\mathcal{A}}^{(2)}$ whose objects are $\mathcal{C}_{\text{Ob}} := \{\mathbb{K}, \mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_1 \oplus \mathbf{X}_2\}$ and whose morphisms are given as all compositions of mappings $\{f_1, f_2, e_{\mathbf{X}_1}, e_{\mathbf{X}_2}\}$ and identity morphisms.

For an n -regular graded algebra $\mathcal{A} = \mathcal{A}^{reg}(\Theta_1, \Theta_2, \dots, \Theta_n)$ generated by a set of regular odd generators $\{\Theta_i; i = 1, \dots, n\}$ we introduce the following decomposition

$$(19) \quad \mathcal{A} = \bigoplus_{i=1}^n \mathbf{X}_i,$$

such that $\mathbf{L}_{\Theta_i}(x) \in \mathbf{X}_{i+1}$ for every $x \in \mathbf{X}_i$, and $i = 1, \dots, n-1$. The space $\mathbf{X}_1$ is the $\mathbb{K}$ -linear span of all monomials in generators ended by Θ_i . For $i = 2, 3, \dots, n$ we define

$$(20) \quad \mathbf{X}_i := \text{lin}_{\mathbb{K}}\{\mathbf{L}_{\Theta_{i-1}} \circ \dots \circ \mathbf{L}_{\Theta_1}(x) : x \in \mathbf{X}_1; i = 2, \dots, n\}.$$

and obtain

$$(21) \quad \begin{aligned} \mathbf{X}_2 &:= \text{lin}_{\mathbb{K}}\{\mathbf{L}_{\Theta_1}(x) : x \in \mathbf{X}_1\}, \\ \mathbf{X}_3 &:= \text{lin}_{\mathbb{K}}\{\mathbf{L}_{\Theta_2} \circ \mathbf{L}_{\Theta_1}(x) : x \in \mathbf{X}_1\}, \\ &\vdots \\ \mathbf{X}_n &:= \text{lin}_{\mathbb{K}}\{\mathbf{L}_{\Theta_{n-1}} \circ \dots \circ \mathbf{L}_{\Theta_1}(x) : x \in \mathbf{X}_1\}. \end{aligned}$$

We use operators $\mathbf{L}_a$ in order to introduce a collection of linear mappings $(\mathbf{X}, f_i) := \{f_i : \mathbf{X}_i \rightarrow \mathbf{X}_{i+1}; i = 1, \dots, n\}$. We define these mappings by the following formula

$$(22) \quad f_i(x) := \mathbf{L}_{\Theta_i}(x) = \Theta_i x$$

for every $x \in \mathbf{X}_i$, $i = 1, \dots, n-1$. One can calculate that we have the higher regularity relation

$$(23) \quad f_1 \circ f_2 \circ \dots \circ f_n \circ f_1 = f_1$$

and corresponding cyclic permutations. In this case there is also a specific category $\mathcal{C}_{\mathcal{A}}^{(n)}$ which contains all spaces and mappings considered above.

6. CATEGORIES AND OBSTRUCTIONS

6.1. Algebras via categories. Recall (see e.g. [22]) that a *category* $\mathcal{C} = (\mathcal{C}_{\text{Ob}}, \mathcal{C}_{\text{Mor}}, \mathbf{c})$ contains

- (i) a collection $\mathcal{C}_{\text{Ob}}$ of objects;
- (ii) a collection $\mathcal{C}_{\text{Mor}}$ of morphisms (arrows)

$$(24) \quad \mathcal{C}_{\text{Mor}} = \bigcup_{X, Y \in \mathcal{C}_{\text{Ob}}} \mathcal{C}(X, Y)$$

- (iii) an associative composition $\mathbf{c}$ of morphisms

$$(25) \quad \mathbf{c} : \mathcal{C}(X, Y) \times \mathcal{C}(Y, \mathcal{W}) \rightarrow \mathcal{C}(X, \mathcal{W})$$

The collection $\mathcal{C}_{\text{Mor}}$ is the union of mutually disjoint sets $\mathcal{C}(X, Y)$ of morphisms $X \rightarrow Y$ from X to Y defined for every pair of objects $X, Y \in \mathcal{C}_{\text{Ob}}$. It may happen that for a pair $X, Y \in \mathcal{C}_{\text{Ob}}$ the set $\mathcal{C}(X, Y)$ is empty.

An *opposite* (or *dual*) category of a category $\mathcal{C} = (\mathcal{C}_{\text{Ob}}, \mathcal{C}_{\text{Mor}}, \mathbf{c})$ is a category $\mathcal{C}^{op} = (\mathcal{C}_{\text{Ob}}, \mathcal{C}_{\text{Mor}}^{op}, \mathbf{c}^{op})$ equipped with the same collection of objects $\mathcal{C}_{\text{Ob}}$ as the category $\mathcal{C}$ but with reversed all arrows

$$(26) \quad \mathcal{C}^{op}(X, Y) \equiv \mathcal{C}(Y, X).$$

and reversed composition $\mathbf{c}^{op}$. If $\mathfrak{D}_{\mathcal{C}}$ is a diagram built from objects and morphisms of the category $\mathcal{C}$, then the same diagram with reversed all arrows $\mathfrak{D}_{\mathcal{C}^{op}}$ is said to be *dual* to $\mathfrak{D}_{\mathcal{C}}$.

Let $\mathcal{C}$ and $\mathcal{D}$ be two categories. A *functor* $\mathfrak{F} : \mathcal{C} \rightarrow \mathcal{D}$ of $\mathcal{C}$ into $\mathcal{D}$ is a pair of maps

$$(27) \quad \mathfrak{F}_{\text{Ob}} : \mathcal{C}_{\text{Ob}} \rightarrow \mathcal{D}_{\text{Ob}}, \quad \mathfrak{F}_{\text{Mor}} : \mathcal{C}_{\text{Mor}} \rightarrow \mathcal{D}_{\text{Mor}}$$

such that

$$(28) \quad \mathfrak{F}_{\text{Mor}}(\varphi \circ \psi) = \mathfrak{F}_{\text{Mor}}(\varphi) \circ \mathfrak{F}_{\text{Mor}}(\psi)$$

for every morphisms $\varphi : Y \rightarrow Z$ and $\psi : X \rightarrow Y$ of $\mathcal{C}$. The generalization to multifunctors is obvious. For instance an m -ary functor $\mathfrak{F} : \mathcal{C}^{\times m} \rightarrow \mathcal{D}$ sends an m -tuple of objects of $\mathcal{C}$ into an object of $\mathcal{D}$.

A *natural transformation* of functors $\mathfrak{s} : \mathfrak{F} \rightarrow \mathfrak{G}$ of $\mathfrak{F}$ into $\mathfrak{G}$ is a collection of morphisms

$$(29) \quad \mathfrak{s} = \{\mathfrak{s}_X : \mathfrak{F}(X) \rightarrow \mathfrak{G}(X), \quad X \in \mathcal{C}_{\text{Ob}}\},$$

such that

$$(30) \quad \mathfrak{s}_Y \circ \mathfrak{F}(\psi) = \mathfrak{G}(\psi) \circ \mathfrak{s}_X$$

for every morphism $\psi : X \rightarrow Y$ of $\mathcal{C}$. The set of all natural transformations of $\mathfrak{F}$ into $\mathfrak{G}$ we denote by $\mathfrak{Nat}(\mathfrak{F}, \mathfrak{G})$. If $\mathfrak{F} \equiv \mathfrak{G}$, then we say that the natural transformation $\mathfrak{s} : \mathfrak{F} \rightarrow \mathfrak{G}$ is a natural transformation of $\mathfrak{F}$ into itself.

We can use functors and natural transformations in order to describe certain algebraic structures in categories. Let $\mathcal{C}$ be a category and let $\mathfrak{F}$ be a functor $\mathfrak{F} : \mathcal{C} \rightarrow \mathcal{C}$. A binary multiplication is a natural transformation $\mathfrak{m} : \mathfrak{F} \times \mathfrak{F} \rightarrow \mathfrak{F}$ satisfying the associativity condition $\mathfrak{m} \circ (\mathfrak{m} \times \mathfrak{F}) = \mathfrak{m} \circ (\mathfrak{F} \times \mathfrak{m})$. Similarly, a comultiplication is a natural transformation $\Delta : \mathfrak{F} \rightarrow \mathfrak{F} \times \mathfrak{F}$ satisfying the corresponding coassociativity condition $(\Delta \times \mathfrak{F}) \circ \Delta = (\mathfrak{F} \times \Delta) \circ \Delta$. A monoidal category $\mathcal{C} \equiv \mathcal{C}(\otimes, \mathbb{K})$ is a category $\mathcal{C}$ equipped with a monoidal operation (a bifunctor) $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$, a unit object $\mathbb{K}$ satisfying some known axioms [18, 28].

We define a (left) $\vee$ -operation

$$(31) \quad X^\vee := \mathcal{C}(X, \mathbb{K})$$

such that we have

$$(32) \quad X^{\vee\vee} = X, \quad (X \otimes Y)^\vee = Y^\vee \otimes X^\vee$$

for every $X, Y \in \mathcal{C}_{\text{Ob}}$. If $\psi : X \rightarrow Y$ is a morphism in $\mathcal{C}$, then there is also a corresponding *dual morphism* $\psi^\vee : Y^\vee \rightarrow X^\vee$

$$(33) \quad \psi^\vee(v^\vee) := u^\vee, \quad u \in X, v \in Y \text{ if and only if } \psi(u) = v, \quad u^\vee \in X^\vee, v^\vee \in Y^\vee.$$

The corresponding category $\mathcal{C}$ is called a monoidal category with duality [18, 28].

A class $\mathbf{g}(\mathcal{C}) := \{\mathbf{g}_X \equiv \langle \dots | \dots \rangle_X : X \in \mathcal{C}_{\text{Ob}}\}$ of $\mathbb{K}$ -valued mappings $\mathbf{g}_X : X^\vee \otimes X \rightarrow \mathbb{K}$ together with a class $\mathbf{h}(\mathcal{C}) := \{\mathbf{h}_X : \mathbb{K} \rightarrow X \otimes X^\vee\}$ such that the following diagrams

$$\begin{array}{ccc} Y^\vee \otimes X^\vee \otimes X \otimes Y & \xrightarrow{\text{id}_{Y^\vee} \otimes \mathbf{g}_X \otimes \text{id}_Y} & Y^\vee \otimes Y \\ \parallel & & \downarrow \mathbf{g}_Y \\ (X \otimes Y)^\vee \otimes (X \otimes Y) & \xrightarrow{\quad \quad \quad} & \mathbb{K} \\ & \mathbf{g}_{X \otimes Y} & \end{array}$$

and

$$\begin{array}{ccc}
 Y^\vee \otimes X \otimes X^\vee & \xrightarrow{\text{id}_{Y^\vee} \otimes \psi \otimes \text{id}_{X^\vee}} & Y^\vee \otimes Y \otimes X^\vee \\
 \text{id}_{Y^\vee} \otimes h_X \uparrow & & \downarrow g_Y \otimes \text{id}_{X^\vee} \\
 Y^\vee & \xrightarrow{\psi^\vee} & X^\vee
 \end{array}$$

is said to be a pairing in $\mathcal{C}$ [28].

6.2. Obstructed categories. Let $\mathcal{C}$ be a category with invertible and noninvertible morphisms [12, 11] and an equivalence. By an equivalence in $\mathcal{C}$ we mean a class of morphisms $\mathcal{C}^P = \bigcup_{X,Y \in \mathcal{C}_{\text{Ob}}} \mathcal{C}^P(X,Y)$, where $\mathcal{C}^P(X,Y)$ is a subset of invertible morphisms in $\mathcal{C}(X,Y)$ satisfying the property P. Two objects X, Y of the category $\mathcal{C}$ are equivalent, if and only if there is an morphism $X \xrightarrow{\psi} Y$ in $\mathcal{C}^P(X,Y)$ such that $\psi^{-1} \circ \psi = \text{id}_X$ and $\psi \circ \psi^{-1} = \text{id}_Y$.

Definition 4. A sequence of (invertible and noninvertible) morphisms

$$(34) \quad X_1 \xrightarrow{\psi_1} X_2 \xrightarrow{\psi_2} \dots \xrightarrow{\psi_{n-1}} X_n \xrightarrow{\psi_n} X_1$$

such that there is an (endo-)morphism $\mathbf{e}_{X_1}^{(n)} : X_1 \longrightarrow X_1$ defined by

$$(35) \quad \mathbf{e}_{X_1}^{(n)} := \psi_n \circ \dots \circ \psi_2 \circ \psi_1$$

and subjects to the following relation

$$(36) \quad \psi_1 \circ \psi_n \circ \dots \circ \psi_2 \circ \psi_1 = \psi_1$$

is said to be a *regular n -cocycle* on $\mathcal{C}$ and it is denoted by (X, ψ) .

Two mappings f_1 and f_2 defined in (16) form a regular 2-cocycle on $\mathcal{C}_{\mathcal{A}}^{(2)}$. Mappings $f_1, \dots, f_n$ from (23) form a regular n -cocycle on $\mathcal{C}_{\mathcal{A}}^{(n)}$.

Definition 5. The morphism $\mathbf{e}_{X_1}^{(n)}$ is said to be an *obstruction* of the object X_1 corresponding to the regular n -cocycle $\psi = (\psi_1, \dots, \psi_n)$ on $\mathcal{C}$. The obstruction (endo-)morphisms $\mathbf{e}_{X_i}^{(n)} : X_i \longrightarrow X_i$ corresponding for $i = 2, \dots, n$ are defined by a suitable cyclic permutation of above sequence.

Definition 6. If the obstruction (endo-)morphisms $\mathbf{e}_X^{(n)} : X \longrightarrow X$ are defined for every object $X \in \mathcal{C}_{\text{Ob}}$, then the mapping $\varepsilon^{(n)} : X \rightarrow \mathbf{e}_X^{(n)} \in \mathcal{C}(X, X)$ is called a *regular n -cocycle obstruction structure* on $\mathcal{C}$.

It is obvious that for an usual category all $\mathbf{e}_X^{(n)}$ are equal to identity $\mathbf{e}_X^{(n)} = \text{Id}_X$. We are interested with categories for which the obstruction $\mathbf{e}_X^{(n)}$ differs from the identity (see e.g. [9]).

Definition 7. A category $\mathcal{C}$ equipped with a regular n -cocycle obstruction structure $\varepsilon^{(n)} : X \in \mathcal{C}_{\text{Ob}} \rightarrow \mathbf{e}_X^{(n)} \in \mathcal{C}(X, X)$ such that $\mathbf{e}_X^{(n)} \neq \text{id}_X$ for some $X \in \mathcal{C}_{\text{Ob}}$ is called an *obstructed category*. The minimum number $n = n_{\text{obstr}}$ for which it occurs will define a quantitative measure of obstruction n_{obstr} .

The categories $\mathcal{C}_{\mathcal{A}}^{(2)}$ and $\mathcal{C}_{\mathcal{A}}^{(n)}$ considered above are obstructed.

Let (Y, φ) a regular n -cocycle, i.e. a sequence of morphisms $Y_1 \xrightarrow{\varphi_1} Y_2 \xrightarrow{\varphi_2} \dots \xrightarrow{\varphi_{n-1}} Y_n \xrightarrow{\varphi_n} Y_1$ such that $\mathbf{e}_{Y_1}^{(n)} := \varphi_n \circ \dots \circ \varphi_2 \circ \varphi_1$.

Definition 8. A sequence of morphisms $\alpha := (\alpha_1, \dots, \alpha_n)$ such that the diagram

$$(37) \quad \begin{array}{ccccccc} X_1 & \xrightarrow{\psi_1} & X_2 & \xrightarrow{\psi_2} \dots \xrightarrow{\psi_{n-1}} & X_n & \xrightarrow{\psi_n} & X_1 \\ \downarrow \alpha_1 & & \downarrow \alpha_2 & & \downarrow \alpha_n & & \downarrow \alpha_1 \\ Y_1 & \xrightarrow{\varphi_1} & Y_2 & \xrightarrow{\varphi_2} \dots \xrightarrow{\varphi_{n-1}} & Y_n & \xrightarrow{\varphi_n} & Y_1 \end{array}$$

is commutative, is said to be a *regular n -cocycle morphism* from (X, ψ) to (Y, φ) and is denoted by $\alpha : (X, \psi) \rightarrow (Y, \varphi)$.

Observe that for a regular n -cocycle morphism $\alpha : (X, \psi) \rightarrow (Y, \varphi)$ we have the relation

$$(38) \quad \alpha_1 \circ \mathbf{e}_{X_1}^{(n)} = \mathbf{e}_{Y_1}^{(n)} \circ \alpha_1.$$

Definition 9. A regular n -cocycle obstruction morphism $\beta : (X, \psi) \rightarrow (X', \varphi)$ which sends the object X_i into equivalent object X'_i and morphism ψ_i into φ_i is said to be *obstruction n -cocycle equivalence*. The corresponding obstructions $\mathbf{e}_X^{(n)}$ and $\mathbf{e}_{X'}^{(n)}$ are also said to be *equivalent*.

Let $\mathcal{C}$ and $\mathcal{D}$ be two obstructed categories. The morphisms $\mathbf{e}_X^{(n)}$ can be used to extend the notion of functors. Let $\mathfrak{F} : \mathcal{C} \rightarrow \mathcal{D}$ be a functor defined as usual as a pair of mappings $(\mathfrak{F}_{\text{Ob}}, \mathfrak{F}_{\text{Mor}})$.

Definition 10. A set of category mappings $\mathfrak{F}^{(n)} : \mathcal{C} \rightarrow \mathcal{D}$ defined as $(\mathfrak{F}_{\text{Ob}}^{(n)}, \mathfrak{F}_{\text{Mor}}^{(n)})$ such that

$$(39) \quad \mathfrak{F}_{\text{Ob}}^{(n)} \equiv \mathfrak{F}_{\text{Ob}}, \quad \mathfrak{F}_{\text{Mor}}^{(n)} \left(\mathbf{e}_{\mathcal{C}_{\text{Ob}}(X)}^{(n)} \right) = \mathbf{e}_{\mathcal{D}_{\text{Ob}}(X)}^{(n)}, \quad \mathfrak{F}_{\text{Mor}}^{(1)} \equiv \mathfrak{F}_{\text{Mor}},$$

where $\mathfrak{F} = (\mathfrak{F}_{\text{Ob}}, \mathfrak{F}_{\text{Mor}})$ is a functor, is said to be *n -regular obstructed functor*.

We see that all the standard definitions of a functor are not changed, but the preservation of identity $\mathfrak{F}_{\text{Mor}}(\text{id}_X) = \text{id}_Y$, where $Y = \mathfrak{F}_{\text{Ob}}(X)$, $X \in \mathcal{C}_{\text{Ob}}$, $Y \in \mathcal{D}_{\text{Ob}}$ is replaced by requirement of preservation of the morphisms $\mathbf{e}_X^{(n)}$, and so the obstructed functor $\mathfrak{F}^{(n)}$ becomes n -dependent.

Lemma 11. *Let the sequence*

$$(40) \quad X_1 \xrightarrow{\psi_1} X_2 \xrightarrow{\psi_2} \dots \xrightarrow{\psi_{n-1}} X_n \xrightarrow{\psi_n} X_1$$

be a regular n -cocycle in the category $\mathcal{C}$. If $\mathfrak{F}^{(n)} : \mathcal{C} \rightarrow \mathcal{D}$ is an n -regular functor, then

$$(41) \quad \mathfrak{F}_{\text{Ob}}^{(n)}(X_1) \xrightarrow{\mathfrak{F}_{\text{Mor}}^{(n)}(\psi_1)} \mathfrak{F}_{\text{Ob}}^{(n)}(X_2) \xrightarrow{\mathfrak{F}_{\text{Mor}}^{(n)}(\psi_2)} \dots \xrightarrow{\mathfrak{F}_{\text{Mor}}^{(n)}(\psi_{n-1})} \mathfrak{F}_{\text{Ob}}^{(n)}(X_n) \xrightarrow{\mathfrak{F}_{\text{Mor}}^{(n)}(\psi_n)} \mathfrak{F}_{\text{Ob}}^{(n)}(X_1)$$

is a regular n -cocycle in the category $\mathcal{D}$, i.e. $Y_1 \xrightarrow{\varphi_1} Y_2 \xrightarrow{\varphi_2} \dots \xrightarrow{\varphi_{n-1}} Y_n \xrightarrow{\varphi_n} Y_1$, $Y_i = \mathfrak{F}_{\text{Ob}}^{(n)}(X_i) \in \mathcal{D}_{\text{Ob}}$, $\varphi_i = \mathfrak{F}_{\text{Mor}}^{(n)}(X_i) \in \mathcal{D}_{\text{Mor}}$.

Proof: Because $\mathfrak{F}_{\text{Ob}}^{(n)} = \mathfrak{F}_{\text{Ob}}$, we need to prove that $\mathfrak{F}_{\text{Mor}}^{(n)}(\psi_i) \circ \mathbf{e}_{X_i}^{(n)} = \mathfrak{F}_{\text{Mor}}^{(n)}(\psi_i)$. Indeed, we have

$$\mathfrak{F}_{\text{Mor}}^{(n)}(\psi_i) = \mathfrak{F}_{\text{Mor}}^{(n)}\left(\psi_i \circ \mathbf{e}_{X_i}^{(n)}\right) = \mathfrak{F}_{\text{Mor}}^{(n)}(\psi_i) \circ \mathfrak{F}_{\text{Mor}}^{(n)}\left(\mathbf{e}_{X_i}^{(n)}\right) = \mathfrak{F}_{\text{Mor}}^{(n)}(\psi_i) \circ \mathbf{e}_{X_i}^{(n)}.$$

□

Multifunctors can be regularized in a similar way. Next we “regularize” natural transformations. Let $\mathfrak{F}^{(n)}$ and $\mathfrak{G}^{(n)}$ be two n -regular functors of the category $\mathcal{C}$ into the category $\mathcal{D}$.

Definition 12. A natural transformation $\mathfrak{s} : \mathfrak{F}^{(n)} \rightarrow \mathfrak{G}^{(n)}$ is a collection of morphisms $\mathfrak{s} = \{\mathfrak{s}_X : \mathfrak{F}_{\text{Ob}}(X) \rightarrow \mathfrak{G}_{\text{Ob}}(X), X \in \mathcal{C}_{\text{Ob}}\}$ such that

$$(42) \quad \mathfrak{s}_Y \circ \mathfrak{F}_{\text{Mor}}^{(n)}(\alpha) = \mathfrak{G}_{\text{Mor}}^{(n)}(\alpha) \circ \mathfrak{s}_X,$$

for every regular morphism $\alpha : X \rightarrow Y$, is said to be *n -regular natural transformation*.

Definition 13. A monoidal category $\mathcal{C} \equiv \mathcal{C}(\otimes, \mathbb{K})$ equipped with a family of obstruction morphisms $\mathbf{E}^{(n)} = \{\mathbf{e}_X^{(n)} : X \in \mathcal{C}_{\text{Ob}}; n = 1, 2, \dots\}$ satisfying the “homomorphism” condition

$$(43) \quad \mathbf{e}_{X \otimes Y}^{(n)} = \mathbf{e}_X^{(n)} \otimes \mathbf{e}_Y^{(n)}.$$

is said to be an *obstructed monoidal category*.

Let $\mathcal{C}$ be a monoidal category with duality [28, 23]. If (X, ψ) a regular n -cocycle on $\mathcal{C}$, then there is a corresponding regular n -cocycle $(X^\vee, \psi^\vee)$, called the dual of (X, ψ) . This means that we have a sequence

$$(44) \quad X_1^\vee \xrightarrow{\psi_n^\vee} X_n^\vee \xrightarrow{\psi_{n-1}^\vee} \dots \xrightarrow{\psi_2^\vee} X_2^\vee \xrightarrow{\psi_1^\vee} X_1^\vee$$

such that there is an (endo-)morphism $\mathbf{e}_{X_1^\vee}^{(n)} : X_1^\vee \rightarrow X_1^\vee$ defined by the following equation

$$(45) \quad \mathbf{e}_{X_1^\vee}^{(n)} := \psi_n^\vee \circ \dots \circ \psi_2^\vee \circ \psi_1^\vee$$

and subjects to the following relation

$$(46) \quad \psi_1^\vee \circ \psi_n^\vee \circ \dots \circ \psi_2^\vee \circ \psi_1^\vee = \psi_1^\vee,$$

where $X_i^\vee = \mathcal{C}(X_i, \mathbb{K})$, $\psi_i^\vee : X_{i+1}^\vee \rightarrow X_i^\vee$, $i = 1, \dots, n-1$. Let $\mathbf{g}_X \equiv \langle \dots | \dots \rangle_X : X^\vee \otimes X \rightarrow \mathbb{K}$ be a pairing in $\mathcal{C}$.

Definition 14. If for every object $X \in \mathcal{C}_{\text{Ob}}$ with an obstruction $\mathbf{e}_X^{(n)}$ there is an object $X^\vee$ with an obstruction $\mathbf{e}_{X^\vee}^{(n)}$ such that

$$(47) \quad \langle \mathbf{e}_{X^\vee}^{(n)}(x^\vee), x \rangle = \langle x^\vee | \mathbf{e}_X^{(n)}(x) \rangle_{\mathcal{U}},$$

for $x^\vee \in X^\vee$, $x \in X$, then the category $\mathcal{C}$ is said to be an *obstructed category with duality*.

The category $\mathcal{C}_{\mathcal{A}}^{(2)}$ above is an obstructed category with duality, where.

$$(48) \quad X_1^\vee := \mathcal{C}_{\mathcal{A}}^{(2)}(X_1, \mathbb{K}), \quad X_2^\vee := \mathcal{C}_{\mathcal{A}}^{(2)}(X_2, \mathbb{K}),$$

are spaces of linear functionals on X_1 and X_2 , respectively, and

$$(49) \quad X_1^\vee := \text{lin}_{\mathbb{K}}\{\Xi_1, \Xi_2\Xi_1\}, \quad X_2^\vee := \text{lin}_{\mathbb{K}}\{\Xi_2, \Xi_1\Xi_2\}.$$

where

$$(50) \quad \langle \Xi_1 | \Theta_1 \rangle_{X_1} := 1, \quad \langle \Xi_2 | \Theta_2 \rangle_{X_2} := 1$$

and

$$(51) \quad \langle \Xi_1\Xi_2 | \Theta_2\Theta_1 \rangle_{X_1} := 1 \quad \langle \Xi_2\Xi_1 | \Theta_1\Theta_2 \rangle_{X_2} := 1.$$

The monoidal product is the $\mathbb{K}$ -tensor product [28, 23].

7. REGULAR ALGEBRAS AND BIALGEBRAS

Let $\mathcal{C}$ be an obstructed monoidal category [14, 15].

Definition 15. An algebra $\mathcal{A}^{reg}$ in the category $\mathcal{C}$ such that the multiplication $m : \mathcal{A}^{reg} \otimes \mathcal{A}^{reg} \rightarrow \mathcal{A}^{reg}$ is a regular n -cocycle morphism

$$(52) \quad m \circ (\mathbf{e}_{\mathcal{A}}^{(n)} \otimes \mathbf{e}_{\mathcal{A}}^{(n)}) = \mathbf{e}_{\mathcal{A}}^{(n)} \circ m,$$

is said to be a *regular n -cocycle algebra*.

Lemma 16. *The graded algebra $\mathcal{A}^{reg}(\Theta_1, \Theta_2)$ is a regular 2-cocycle algebra in the category $\mathcal{C}_{\mathcal{A}}$.*

Proof: Let $\mathbf{a}, \mathbf{b} \in \mathcal{A}^{reg}(\Theta_1, \Theta_2)$ have the form (8). Then for the obstruction $\mathbf{e}_{\mathcal{A}}^{(2)}$ we calculate

$$(53) \quad \mathbf{e}_{\mathcal{A}}^{(2)}(\mathbf{a}) := 1 + a_2\Theta_1 + a_1\Theta_2 + a_{21}\Theta_1\Theta_2 + a_{12}\Theta_2\Theta_1$$

So the condition (52) for $\mathbf{e}_{\mathcal{A}}^{(2)}(\mathbf{a}), \mathbf{e}_{\mathcal{A}}^{(2)}(\mathbf{b}), \mathbf{e}_{\mathcal{A}}^{(2)}(\mathbf{c})$ in components is

$$\begin{aligned} c_1 &= a_1 + b_1 + a_2b_{12} + a_{21}b_2, \\ c_2 &= a_2 + b_2 + a_1b_{21} + a_{12}b_1, \\ c_{12} &= a_{12} + b_{12} + a_1b_2 + a_{12}b_{12}, \\ c_{21} &= a_{21} + b_{21} + a_2b_1 + a_{21}b_{21}. \end{aligned}$$

□

Note that in case of ground field is complex among obstructions $\mathbf{e}_{\mathcal{A}}^{(2)}(\mathbf{a})$ there are exist only two ones which are idempotent as elements, i. e. satisfy the equation $\mathbf{e}_{\mathcal{A}}^{(2)}(\mathbf{a})\mathbf{e}_{\mathcal{A}}^{(2)}(\mathbf{a}) = \mathbf{e}_{\mathcal{A}}^{(2)}(\mathbf{a})$, and they do not depend from $\mathbf{a}$

$$\mathbf{e}_{\mathcal{A}}^{(2)\text{idemp}} = 1 + \Theta_1 + \Theta_2 + \left(-\frac{1}{2} \pm i\frac{\sqrt{3}}{2}\right)\Theta_1\Theta_2 + \left(-\frac{1}{2} \mp i\frac{\sqrt{3}}{2}\right)\Theta_2\Theta_1.$$

There is a graded algebra $\mathcal{A}^{reg, \vee}(\Xi_1, \Xi_2)$ generated by Ξ_1 and Ξ_2 subject to the relations

$$(54) \quad \begin{aligned} \Xi_1^2 &= \Xi_2^2 = 0, \\ \Xi_1\Xi_2\Xi_2 &= \Xi_1, \quad \Xi_2\Xi_1\Xi_2 = \Xi_2. \end{aligned}$$

which is also is a regular 2-cocycle algebra in the category $\mathcal{C}$.

□

Definition 17. A coalgebra $\mathcal{A}^{reg}$ in the category $\mathcal{C}$ such that the comultiplication $\Delta_{reg} : \mathcal{A}^{reg} \longrightarrow \mathcal{A}^{reg} \otimes \mathcal{A}^{reg}$ is a regular n -cocycle morphism

$$(55) \quad \Delta_{reg} \circ \mathbf{e}_{\mathcal{A}}^{(n)} = (\mathbf{e}_{\mathcal{A}}^{(n)} \otimes \mathbf{e}_{\mathcal{A}}^{(n)}) \circ \Delta_{reg}.$$

is said to be a *regular n -cocycle coalgebra*.

Lemma 18. Let $\mathcal{A}$ be an algebra in an obstructed category with duality, and let $\mathcal{A}^\vee$ be its dual coalgebra

$$(56) \quad \langle \Delta_{reg}(\mathbf{a}^\vee), \mathbf{a}_1 \otimes \mathbf{a}_2 \rangle = \langle \mathbf{a}^\vee, m(\mathbf{a}_1 \otimes \mathbf{a}_2) \rangle,$$

where $\mathbf{a}_1, \mathbf{a}_2 \in \mathcal{A}$, $\mathbf{a}^\vee \in \mathcal{A}^\vee$. If $\mathcal{A}$ is a regular n -cocycle algebra, then $\mathcal{A}^\vee$ is a regular n -cocycle coalgebra.

Proof: Let us apply the regularity condition (52) to the above duality condition (56). Then the lemma follows from relations (43), (47), and (55). $\square$

The graded algebra $\mathcal{A}^\vee(\Xi_1, \Xi_2)$ is also a regular 2-cocycle coalgebra which is dual to $\mathcal{A}^{reg}(\Theta_1, \Theta_2)$ in the category $\mathcal{C}$.

Definition 19. Let $\mathcal{A}^{reg}$ be a regular n -cocycle algebra. If $\mathcal{A}^{reg}$ is also regular coalgebra such that $\Delta_{reg}(\mathbf{ab}) = \Delta_{reg}(\mathbf{a}) \Delta_{reg}(\mathbf{b})$, then it is said to be a *regular n -cocycle almost bialgebra*.

One can also consider the corresponding generalization of Hopf algebras [25, 19].

The graded algebra $\mathcal{A}^{reg}(\Theta_1, \Theta_2)$ is a regular 2-cocycle almost bialgebra, where

$$(57) \quad \Delta_{reg}(\Theta_1) = \Theta_1 \otimes \mathbf{e}_{\mathbf{x}_1} + \mathbf{e}_{\mathbf{x}_1} \otimes \Theta_1,$$

$$(58) \quad \Delta_{reg}(\Theta_2) = \Theta_2 \otimes \mathbf{e}_{\mathbf{x}_2} + \mathbf{e}_{\mathbf{x}_2} \otimes \Theta_2,$$

Similarly, the graded algebra $\mathcal{A}^\vee(\Xi_1, \Xi_2)$ is a regular 2-cocycle almost bialgebra.

Definition 20. Let M^{reg} be an $\mathcal{A}^{reg}$ -module equipped with the $\mathcal{A}^{reg}$ -module action $\rho_M^{reg} : \mathcal{A}^{reg} \otimes M^{reg} \longrightarrow M^{reg}$, where $\mathcal{A}^{reg}$ is a n -regular cocycle algebra. Then M^{reg} is said to be an *n -regular cocycle $\mathcal{A}^{reg}$ -module*, if and only if

$$(59) \quad \rho_M^{reg} \circ (\mathbf{e}_{\mathcal{A}}^{(n)} \otimes \mathbf{e}_M^{(n)}) = \mathbf{e}_M^{(n)} \circ \rho_M^{reg}.$$

8. REGULAR WICK ALGEBRAS

A pair of algebras $\mathcal{A}$ and $\mathcal{A}^\dagger$ equipped with an antilinear and involutive anti-isomorphism $(-)^{\dagger} : \mathcal{A} \rightarrow \mathcal{A}^\dagger$ such that

$$(60) \quad m_{\mathcal{A}^\dagger}(\mathbf{b}^\dagger \otimes \mathbf{a}^\dagger) = (m_{\mathcal{A}}(\mathbf{a} \otimes \mathbf{b}))^\dagger, \quad (\mathbf{a}^\dagger)^\dagger = \mathbf{a},$$

where $\mathbf{a}, \mathbf{b} \in \mathcal{A}$ and $\mathbf{a}^\dagger, \mathbf{b}^\dagger \in \mathcal{A}^\dagger$ are their images under the isomorphism $(-)^{\dagger}$, is said to be the *conjugated* one to the other [6].

Definition 21. Let $(\mathcal{A}, \mathcal{A}^\dagger)$ be a pair of conjugated algebras. A linear mapping $\Psi : \mathcal{A}^\dagger \otimes \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}^\dagger$ such that

$$(61) \quad \begin{aligned} \Psi \circ (\text{id}_{\mathcal{A}^\dagger} \otimes m_{\mathcal{A}}) &= (m_{\mathcal{A}} \otimes \text{id}_{\mathcal{A}^\dagger}) \circ (\text{id}_{\mathcal{A}} \otimes \Psi) \circ (\Psi \otimes \text{id}_{\mathcal{A}}), \\ \Psi \circ (m_{\mathcal{A}^\dagger} \otimes \text{id}_{\mathcal{A}}) &= (\text{id}_{\mathcal{A}} \otimes m_{\mathcal{A}^\dagger}) \circ (\Psi \otimes \text{id}_{\mathcal{A}^\dagger}) \circ (\text{id}_{\mathcal{A}^\dagger} \otimes \Psi) \end{aligned}$$

is said to be a *cross symmetry* or *generalized twist* [6].

Definition 22. The tensor product $\mathcal{A} \otimes \mathcal{A}^\dagger$ equipped with the multiplication

$$(62) \quad m_\Psi := (m_{\mathcal{A}} \otimes m_{\mathcal{A}^\dagger}) \circ (\text{id}_{\mathcal{A}} \otimes \Psi \otimes \text{id}_{\mathcal{A}^\dagger})$$

is an associative algebra called a *Hermitian Wick algebra* [6, 17] and it is denoted by $\mathcal{W} = \mathcal{W}_\Psi(\mathcal{A}) = \mathcal{A} \rtimes_\Psi \mathcal{A}^\dagger$.

Definition 23. Let $(\mathcal{A}, \mathcal{A}^\dagger)$ be a pair of conjugated algebras, and let $\Psi : \mathcal{A}^\dagger \otimes \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}^\dagger$ be a cross symmetry. If both algebras $\mathcal{A}$ and $\mathcal{A}^\dagger$ are n -regular cocycle algebras and

$$(63) \quad (\mathbf{e}_{\mathcal{A}}^{(n)} \otimes \mathbf{e}_{\mathcal{A}^\dagger}^{(n)}) \circ \Psi^{reg} := \Psi^{reg} \circ (\mathbf{e}_{\mathcal{A}^\dagger}^{(n)} \otimes \mathbf{e}_{\mathcal{A}}^{(n)}),$$

then the mapping Ψ^{reg} is said to be *n -regular cocycle cross symmetry* and the corresponding algebra $\mathcal{W}^{reg} = \mathcal{W}_{\Psi^{reg}}(\mathcal{A}) = \mathcal{A} \rtimes_{\Psi^{reg}} \mathcal{A}^\dagger$ is called *n -regular cocycle Wick algebra*.

A natural regular generalization of (62) is given by

Definition 24. The tensor product $\mathcal{A} \otimes \mathcal{A}^\dagger = \mathcal{A}^{reg} \otimes \mathcal{A}^{reg, \dagger}$ equipped with the multiplication

$$(64) \quad m_\Psi^{reg} := (m_{\mathcal{A}} \otimes m_{\mathcal{A}^\dagger}) \circ (\mathbf{e}_{\mathcal{A}}^{(n)} \otimes \Psi^{reg} \otimes \mathbf{e}_{\mathcal{A}^\dagger}^{(n)})$$

is an associative algebra called a *regular Hermitian Wick algebra* and it is denoted by $\mathcal{W}^{reg} = \mathcal{W}_{\Psi^{reg}}(\mathcal{A}) = \mathcal{A} \rtimes_{\Psi^{reg}} \mathcal{A}^\dagger$.

Let us describe the regular Wick algebra corresponding to the algebra $\mathcal{A}^{reg}(\Theta_1, \Theta_2)$. In this case $\mathcal{A}^\dagger = \mathcal{A}^{reg, \vee}(\Xi_1, \Xi_2)$. Let us assume that

$$(65) \quad \begin{aligned} \Psi^{reg}(\Xi_1 \otimes \Theta_1) &= e_{\mathbf{x}_1} - \Theta_1 \otimes \Xi_1 & \Psi^{reg}(\Xi_1 \otimes \Theta_2) &= \Theta_2 \otimes \Xi_1, \\ \Psi^{reg}(\Xi_2 \otimes \Theta_2) &= e_{\mathbf{x}_2} - \Theta_2 \otimes \Xi_2 & \Psi^{reg}(\Xi_2 \otimes \Theta_1) &= \Theta_1 \otimes \Xi_2, \end{aligned}$$

where $e_{\mathbf{x}_i}$ are defined in (17). We use the (61) relations in order to calculate $\Psi^{reg}(\Xi_1 \otimes \Theta_1 \otimes \Theta_2)$ and others similar expressions. We obtain

$$(66) \quad \begin{aligned} \Psi^{reg}(\Xi_1 \otimes \Theta_1 \otimes \Theta_2) &= \Psi^{reg}(\Psi^{reg}(\Xi_1 \otimes \Theta_1) \otimes \Theta_2) \\ &= \Theta_2 - \Theta_1 \Psi^{reg}(\Xi_1 \otimes \Theta_2) = \Theta_2 - \Theta_1 \Theta_2 \otimes \Xi_1 \end{aligned}$$

Thus using (65) and (66) we can construct an n -regular cocycle Wick algebra $\mathcal{W}^{reg} = \mathcal{W}_{\Psi^{reg}}(\mathcal{A}^{reg}(\Theta_1, \Theta_2))$.

Acknowledgements. S.D. is grateful to Alexander Stolin and Leonid Vaksman for useful discussions and to Jerzy Lukierski for kind hospitality at the University of Wroclaw, where this work was finished. The work of W.M. is partially sponsored by Polish Committee for Scientific Research (KBN) under Grant No 5P03B05620.

REFERENCES

- [1] J. C. Baez, *An introduction to n -categories*, Univ. California preprint, **q-alg/9705009**, Riverside, 1997.
- [2] J. C. Baez and J. Dolan, *Higher-dimensional algebra and topological quantum field theory*, J. Math. Phys. **36** (1995), 6073–6105.
- [3] ———, *Categorification*, Univ. California preprint, **math/9802029**, Riverside, 1998.
- [4] ———, *From finite sets to Feynman diagrams*, Univ. California preprint, **math.QA/0004133**, Riverside, 2000.
- [5] F. A. Berezin, *Introduction to Superanalysis*, Reidel, Dordrecht, 1987.
- [6] A. Borowiec and W. Marcinek, *On crossed product of algebras*, J. Math. Phys. **41** (2000), 6959–6968.

- [7] L. Crane and D. Yetter, *On algebraic structures implicit in topological quantum field theories*, Kansas State Univ. preprint, **hep-th/9412025**, Manhattan, 1994.
- [8] S. Duplij, *On semi-supermanifolds*, Pure Math. Appl. **9** (1998), 283–310.
- [9] ———, *Semisupermanifolds and semigroups*, Krok, Kharkov, 2000.
- [10] S. Duplij and F. Li, *On regular solutions of quantum Yang-Baxter equation and weak Hopf algebras*, Journal of Kharkov National University, ser. Nuclei, Particles and Fields **521** (2001), 15–30.
- [11] S. Duplij and W. Marcinek, *Higher regularity properties of mappings and morphisms*, Univ. Wrocław preprint, IFT UWr 931/00, **math-ph/0005033**, Wrocław, 2000.
- [12] ———, *On higher regularity and monoidal categories*, Kharkov State University Journal (Vestnik KSU), ser. Nuclei, Particles and Fields **481** (2000), 27–30.
- [13] ———, *Semisupermanifolds and regularization of categories*, in Supersymmetric Structures in Mathematics and Physics, (S. Duplij and V. G. Zima, eds.), UkrINTI, Kiev, 2000, pp. 102–115.
- [14] ———, *Noninvertibility, semisupermanifolds and categories regularization*, in Noncommutative Structures in Mathematics and Physics, (S. Duplij and J. Wess, eds.), Kluwer, Dordrecht, 2001, pp. 125–140.
- [15] ———, *Semisupermanifolds and regularization of categories, modules, algebras and Yang-Baxter equation*, in Supersymmetry and Quantum Field Theory, Elsevier Science Publishers, Amsterdam, 2001, pp. 110–115.
- [16] E. Graczynska and Z. Oziewicz, *Hyperequational theories via categories*, Miscellanea Algebraicae **5** (2001), 54–63.
- [17] P. E. T. Jorgensen, L. M. Schmitt, and R. F. Werner, *Positive representation of general commutation relations allowing Wick ordering*, J. Funct. Anal. **134** (1995), 33–45.
- [18] A. Joyal and R. Street, *Braided monoidal categories*, Macquarie University preprint, Mathematics Reports 86008, North Ryde, New South Wales, 1986.
- [19] F. Li, *Weak Hopf algebras and new solutions of Yang-Baxter equation*, J. Algebra **208** (1998), 72–100.
- [20] F. Li and S. Duplij, *Weak Hopf algebras and singular solutions of quantum Yang-Baxter equation*, Zhejiang Univ. preprint, **math.QA/0105064**, Hangzhou, 2001.
- [21] V. V. Lyubashenko, *Squared Hopf Algebras*, Vol. 142 of *Memoirs of AMS*, American Math. Society, Providence, 1999.
- [22] S. MacLane, *Categories for the Working Mathematician*, Springer-Verlag, Berlin, 1971.
- [23] S. Majid, *Quasitriangular Hopf algebras and Yang-Baxter equations*, Int. J. Mod. Phys. **A5** (1990), 1–91.
- [24] D. Nikshych and L. Vainerman, *Finite quantum groupoids and their applications*, Univ. California preprint, **math.QA/0006057**, Los Angeles, 2000.
- [25] F. Nill, *Axioms for weak bialgebras*, Inst. Theor. Phys. FU preprint, **math.QA/9805104**, Berlin, 1998.
- [26] M. Petrich, *Lectures in Semigroups*, Academic Press, Berlin, 1977.
- [27] L. N. Shevrin and A. Y. Ovsyannikov, *Semigroups and Their Subsemigroup Lattices*, Kluwer Academic, Dordrecht, 1996.
- [28] D. N. Yetter, *Quantum groups and representations of monoidal categories*, Math. Proc. Camb. Phil. Soc. **108** (1990), 261–290.

KHARKOV NATIONAL UNIVERSITY, SVOBODA SQ. 4, KHARKOV 61077, UKRAINE
E-mail address: Steven.A.Duplij@univer.kharkov.ua
URL: <http://gluon.physik.uni-kl.de/~duplij>

INSTITUTE OF THEORETICAL PHYSICS, UNIVERSITY OF WROCLAW., PL. MAXA BORNA 9, 50-204
 WROCLAW, POLAND
E-mail address: wmar@ift.univ.wroc.pl